



# Physics

XI & XII

FORMULA SHEET

MADE WITH LOVE

AISHIK GHORAI

for the pals of 12 B and C



AG



# PHYSICAL CONSTANTS

- Speed of Light  $c = 3 \times 10^8 \text{ m/s}$
- Planck constant  $h = 6.63 \times 10^{-34} \text{ Js}$   $hc = 1242 \text{ eV-nm}$
- Gravitation constant  $G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$
- Boltzmann constant  $k = 1.38 \times 10^{-23} \text{ J/K}$
- Molar gas constant  $R = 8.314 \text{ J/mol K}$
- Avogadro's number  $N_A = 6.023 \times 10^{23} / \text{mol}$
- Charge of electron  $e = 1.602 \times 10^{-19} \text{ C}$
- Permeability of vacuum  $\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2$
- Permittivity of vacuum  $\epsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$
- Coulomb constant  $1/4\pi\epsilon_0 = 9 \times 10^9 \text{ N m}^2/\text{C}^2$
- Faraday constant  $F = 96485 \text{ C/mol}$
- Mass of electron  $m_e = 9.1 \times 10^{-31} \text{ kg}$
- Mass of proton  $m_p = 1.6726 \times 10^{-27} \text{ kg}$
- Mass of neutron  $m_n = 1.6744 \times 10^{-27} \text{ kg}$
- Atomic mass unit  $u = 1.66 \times 10^{-27} \text{ kg}$
- Stefan-Boltzmann constant  $\sigma = 5.67 \times 10^{-8} \text{ W/m}^2\text{K}^4$
- Rydberg constant  $R_\infty = 1.097 \times 10^7 / \text{m}$
- Bohr magnetron  $\mu_B = 9.27 \times 10^{-24} \text{ J/T}$
- Bohr radius  $a_0 = 0.529 \times 10^{-10} \text{ m}$
- Standard atmosphere  $atm = 1.01325 \times 10^5 \text{ Pa}$
- Wien displacement constant  $b = 2.9 \times 10^{-3} \text{ mK}$



# LAWS OF MOTION

- 1<sup>st</sup> LAW: INERTIA 2<sup>nd</sup> LAW:  $F = d\vec{p}/dt = ma$  3<sup>rd</sup> LAW: Action  $\Rightarrow$  Reaction
- Friction:  $f_{\text{static, maximum}} = \mu_s N$   $f_{\text{kinetic}} = \mu_k N$
- Centripetal force  $= \frac{mv^2}{r} = m\omega^2 r$



**CURVED BANKING**

$$\frac{v^2 \tan \theta}{rg} = \frac{v^2}{1 \mp \mu \tan \theta}$$

# WORK, POWER & ENERGY

- WORK  $= \vec{F} \cdot \vec{s} = F s \cos \theta = \int \vec{F} \cdot d\vec{s}$
- KE  $= \frac{1}{2} m v^2$  (K)
- POTENTIAL ENERGY (U)
- $U_g = mgh$   $\vec{F} = -\frac{dU}{dx}$  (FOR CONSERVATIVE FORCES)
- $U_{\text{spring}} = \frac{1}{2} k x^2$
- $K + U = \text{Conserved}$
- WORK-ENERGY THEOREM:  $W_{\text{net}} = \Delta K$
- POWER  $= dW/dt = \vec{F} \cdot \vec{v}$

# CENTER OF MASS

- $x_{\text{cm}} = \frac{\sum x_i m_i}{\sum m_i} = \frac{\int x dm}{\int dm}$
- $\vec{v}_{\text{cm}} = \frac{\sum m_i \vec{v}_i}{\sum m_i}$   $\vec{F} = m \vec{a}_{\text{cm}}$
- HOLLOW CONE  $= h/3$
- SOLID CONE  $= h/4$
- HOLLOW SPHERE  $= R/2$
- SOLID SPHERE  $= 3R/8$

# VECTORS

- $\vec{a} = a_x \hat{i} + a_y \hat{j} + a_z \hat{k}$   $|\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2}$
- Dot Product  $\vec{a} \cdot \vec{b} = a_x b_x + a_y b_y + a_z b_z = ab \cos \theta$
- Cross Product  $\vec{a} \times \vec{b} = ab \sin \theta$  (AREA)
- $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix} = (a_y b_z - b_y a_z) \hat{i} + (a_z b_x - b_z a_x) \hat{j} + (a_x b_y - b_x a_y) \hat{k}$

# KINEMATICS

- $\vec{v}_{\text{avg}} = \Delta \vec{s} / \Delta t$   $\vec{v}_{\text{inst}} = d\vec{s} / dt$
- $\vec{a}_{\text{avg}} = \Delta \vec{v} / \Delta t$   $\vec{a}_{\text{inst}} = d\vec{v} / dt$
- $s = ut + \frac{1}{2} a t^2$  **RELATIVE VELOCITY**
- $v = u + at$   $\rightarrow v_{A/B} = v_A - v_B$
- $v^2 = u^2 + 2as$






# PROJECTILE MOTION

- $u_x = u \cos \theta$   $u_y = u \sin \theta$
- Time of Flight  $= 2u_y/g \Rightarrow T = 2u \sin \theta / g$
- Range  $= u_x T \Rightarrow R = \frac{u^2 \sin 2\theta}{g}$
- $y = \tan \theta \cdot x - \left( \frac{g}{2u^2 \cos^2 \theta} \right) \cdot x^2$
- Height  $H = \frac{u^2 \sin^2 \theta}{2g}$

# RIGID BODY DYNAMICS

- $\omega = \frac{\Delta \theta}{\Delta t} = \frac{d\theta}{dt}$   $\alpha = \frac{\Delta \omega}{\Delta t} = \frac{d\omega}{dt}$   $\vec{v} = \vec{\omega} \times \vec{r}$   $\vec{a}_{\text{tan}} = \vec{r} \times \vec{\alpha}$   $\vec{a}_{\text{centri}} = \omega^2 r$
- $\vec{L} = \vec{r} \times \vec{p} = m v r_{\perp}$   $\vec{z} = I \alpha = d\vec{L} / dt$
- $\vec{\tau} = \vec{r} \times \vec{F} = r_{\perp} F = r F \sin \theta$
- EQUILIBRIUM:  $F_{\text{net}} = 0 = \vec{z}_{\text{net}}$   $\omega = 2\pi f$   $T = 1/f$   $\omega = v_{\perp} / r$

# MOMENT OF INERTIA



$$I = \sum m_i r_i^2$$



$$I = \int r^2 dm$$

$$R_{\text{GYRATION}} \quad mk^2 = I$$

## KINETIC ENERGY

$$K = \frac{1}{2} m v_c^2 + \frac{1}{2} I_c \omega^2$$

$$K = \frac{1}{2} I_H \omega^2 \quad \left\{ \begin{array}{l} \text{About Hinge} \\ \text{or } I_{OR} \end{array} \right\}$$




## ROLLING MOTION

$$v = \omega r \quad (\text{no slip condition})$$

$$I_{OR} \text{ INSTANTANEOUS AXIS OF ROTATION}$$


$$\vec{v} = \vec{\omega} \times \vec{r}$$

## AXIS THEOREMS



### PERPENDICULAR

$$I_z = I_x + I_y$$



### PARALLEL

$$I_{||} = I_{cm} + md^2$$

# GRAVITATION

$$F = G \frac{Mm}{R^2} \quad \text{POT. ENERGY } (U) = -GMm/R$$


$$g = G \frac{M}{R^2} \quad g' = g \left[ 1 - \frac{d}{R_e} \right] \quad g'' \approx g \left[ 1 - \frac{2h}{R_e} \right]$$



$$V_{\text{ORBITAL}} = \sqrt{GM/R}$$

$$V_{\text{ESCAPE}} = \sqrt{2GM/R}$$

$$g' = g - \omega^2 R_e \cos^2 \theta$$



## KEPLER'S LAWS

- 1<sup>st</sup> Elliptical Orbits, Sun @ foci
- 2<sup>nd</sup> Equal Area in Equal time (A^2)
- 3<sup>rd</sup> T^2 proportional to a^3 (semi major axis)

# SHS




## HOOKE'S LAW F = -kx

$$x = A \sin(\omega t + \phi)$$

$$v = A \omega \cos(\omega t + \phi)$$

$$a = -\omega^2 x = -k/m x$$


$$T = \frac{2\pi}{\omega} = 2\pi \sqrt{m/k}$$



$$K = \frac{1}{2} m v^2$$

$$U = \frac{1}{2} k x^2$$

$$E = K + U = \frac{1}{2} k A^2 = \frac{1}{2} m \omega^2 A^2$$



$$z = k \theta$$

$$T = 2\pi \sqrt{\frac{I}{k}}$$


$$x_1 = A_1 \sin(\omega t)$$

$$x_2 = A_2 \sin(\omega t + \phi)$$

$$x = x_1 + x_2 = A \sin(\omega t + \epsilon)$$

$$A = \sqrt{A_1^2 + A_2^2 + 2A_1 A_2 \cos \phi}$$

$$\tan \epsilon = \frac{A_2 \sin \phi}{A_1 + A_2 \cos \phi}$$



## SERIES

$$\frac{1}{K_{eq}} = \frac{1}{K_1} + \frac{1}{K_2}$$

## PARALLEL

$$K_{eq} = K_1 + K_2$$

# PROPERTIES OF MATTER

$$\text{YOUNG'S MODULUS } (Y) = \frac{F/A}{\Delta l/l}$$

$$\text{SHEAR MODULUS } (\eta) = \frac{F/A}{\tan \theta}$$

$$\text{BULK MODULUS } (B) = -V \frac{\Delta P}{\Delta V}$$

$$\text{COMPRESSIBILITY } (K) = \frac{1}{B} = -\frac{1}{V} \frac{\Delta V}{\Delta P}$$


$$\text{POISSON'S RATIO } (\sigma) = \frac{\text{LATERAL STRAIN}}{\text{LONGITUDINAL STRAIN}} = \frac{\Delta D/D}{\Delta l/l}$$

$$\text{ELASTIC ENERGY } (U) = \frac{1}{2} \text{STRESS} \times \text{STRAIN} \times \text{VOLUME}$$

$$\text{SURFACE TENSION } (S) = F/l$$

$$\text{SURFACE ENERGY } (U) = S \cdot \text{AREA}$$

$$P_{\text{EXCESS}} = \Delta P_{\text{AIR}} = \frac{2S}{R} \quad \Delta P_{\text{SOAP}} = \frac{4S}{R}$$



$$P_{\text{HYDROSTATIC}} = \rho g h \quad F_{\text{BOUOYANT}} = \rho g V$$

$$\text{CONTINUITY } A_1 v_1 = A_2 v_2$$

$$\text{BERNOULLI'S } P + \rho g h + \frac{1}{2} \rho v^2 = \text{Const}$$



$$F_{\text{VISCOUS}} = -\eta A \frac{dv}{dx}$$

$$\text{TORRICELLI'S } v_{\text{EFFLUX}} = \sqrt{2gh}$$

$$\text{STOKE'S LAW } F = 6\pi \eta r v$$

$$v_{\text{TERMINAL}} = \frac{2r^2(\rho - \sigma)g}{9\eta}$$

$$\text{POISEUILLI'S EQN } \frac{\text{VOLUME FLOW}}{\Delta t} = \frac{\pi r^4 \Delta P}{8\eta L}$$



# WAVES

$$\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 y}{\partial t^2}$$


$$Y = A \sin(kx - \omega t) = A \sin\left[2\pi\left(\frac{x}{\lambda} - \frac{t}{T}\right)\right]$$

$$T = \frac{1}{\nu} = \frac{2\pi}{\omega} \quad v = \nu\lambda \quad \text{WAVE NUMBER } (k) = \frac{2\pi}{\lambda}$$

$Y_1 = A_1 \sin(kx - \omega t) \quad Y_2 = A_2 \sin(kx - \omega t + \phi)$   
 $Y = A \sin(kx - \omega t + \epsilon) \quad A^2 = \sqrt{(A_1 + A_2 \cos \phi)^2 + (A_2 \sin \phi)^2}$   
 $\phi = 2n\pi$  (even) : **constructive**  
 $= (2n+1)\pi$  (odd) : **destructive**  
 $\tan \epsilon = \frac{A_2 \sin \phi}{A_1 + A_2 \cos \phi}$   
 $P_{\text{avg}} = 2\pi^2 \mu \nu A v^2 \quad v = \sqrt{\frac{T}{\mu}}$

## STANDING WAVES

$y_1 = A \sin(kx - \omega t) \quad y_2 = A \sin(kx + \omega t)$   
 $Y = 2A \cos kx \sin \omega t$  Node if 0 is zero  $\rightarrow x = (n + \frac{1}{2})\lambda$



## SOUND WAVES

$S = S_0 \sin[\omega(t - x/v)]$   
 $P = P_0 \cos[\omega(t - x/v)]$   
 $P_0 = \left[\frac{\rho c \omega}{v}\right] S_0$   
 $I = \frac{2\pi^2 B}{v} S_0^2 v^2 = \frac{P_0^2 v}{2B} = \frac{P_0}{2\rho v}$

## STANDING LONGITUDINAL WAVES

$P_1 = P_0 \sin[\omega(t - x/v)] \quad P_2 = P_0 \sin[\omega(t + x/v)]$   
 $P = P_1 + P_2 = 2P_0 \cos kx \sin \omega t$   
**CLOSED ORGAN PIPE**  
 $L = (2n+1) \frac{\lambda}{4} \quad v = (2n+1) \frac{v}{4L}$   
**OPEN ORGAN PIPE**  
 $L = n \frac{\lambda}{2} \quad v = n \frac{v}{2L}$

## RESONANCE COLUMN

$L_1 + d = \frac{\lambda}{2} \quad L_2 + d = \frac{3\lambda}{2}$   
 $v = 2(L_2 - L_1) \nu$   
**BEATS** (if  $\omega_1 \approx \omega_2$ )  
 $P_1 = P_0 \sin \omega_1(t - x/v) \quad P_2 = P_0 \sin \omega_2(t - x/v)$   
 $P = 2P_0 \cos \Delta\omega(t - x/v) \sin \omega(t - x/v)$   
 $\omega = \frac{(\omega_1 + \omega_2)}{2}$  Beats  $\rightarrow \Delta\omega = \omega_1 - \omega_2$   
**DOPPLER**  $v = \frac{v + V_o}{v - V_s} v_o$

## LIGHT WAVES

**PLANE WAVES**  $E = E_0 \sin \omega(t - x/v); I = I_0$   
**SPHERICAL WAVES**  $E = \frac{A E_0}{r} \sin \omega(t - r/v); I = \frac{I_0}{r^2}$

**DIFFRACTION**  
 $\Delta x = b \sin \theta \approx b\theta$   
**Minima**  $b\theta = n\lambda$   
**Resolution**  $\sin \theta = \frac{1.22\lambda}{b}$   
 $\theta \sim \tan \theta = y/D$

## YOUNG'S DOUBLE SLIT EXPERIMENT

**Path diff:**  $\Delta x = y \frac{d}{D}$  **Phase diff:**  $\delta = \frac{2\pi}{\lambda} \Delta x$   
**CONSTRUCTIVE**  $\delta = 2n\pi; \Delta x = n\lambda$  **DESTRUCTIVE**  $\delta = (2n+1)\pi; \Delta x = (n + \frac{1}{2})\lambda$   
**Intensity**  $I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \delta$   $I_{\text{max/min}} = (\sqrt{I_1} \pm \sqrt{I_2})^2$   
**Fringe Width**  $w = \lambda \frac{D}{d}$  **Optical Path**  $\Delta x' = \mu \Delta x$   
**LAW OF MALUS**  $I = I_0 \cos^2 \theta$   
**INTERFERENCE THROUGH THIN FILM**  
 $\Delta x = 2\mu d = n\lambda \rightarrow$  **Constructive**  
 $(2n+1)\lambda/2 \rightarrow$  **destructive**

## OPTICS

**REFLECTION**  
 (i)  $\angle i = \angle r$   
 (ii)  $i, r$  & normal in same plane  
 $f = R/2$   
 $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$   
**Magnification**  $m = -\frac{v}{u}$

**REFRACTION**  
 $\mu = \frac{c}{v} = \frac{\text{vacuum}}{\text{medium}}$   
**SNELL'S LAW**  $\mu_1 \sin i = \mu_2 \sin r$   
**APPARENT DEPTH**  $d' = d/\mu$   
**TIR CRITICAL ANGLE**  
 $\mu \sin \theta_c = \sin 90^\circ$

**PRISM**  
 $S = i + i' - A$   
 $\mu = \frac{\sin\left(\frac{A + \delta_{\min}}{2}\right)}{\sin\left(\frac{A}{2}\right)}$   
 $\delta_{\min} = (\mu - 1)A$   
 For small 'A'


**SPHERICAL SURFACE**  
 $\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$   
 $m = \frac{\mu_1 v}{\mu_2 u}$   
**LENS MAKER'S**  $\frac{1}{f} = (\mu - 1)\left(\frac{1}{R_1} - \frac{1}{R_2}\right)$   
**LENS FORMULA**  $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}; m = \frac{v}{u}$   
**POWER**  $P = \frac{1}{f}$   
**THIN LENSES**  $\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} - \frac{d}{f_1 f_2}$

**MICROSCOPE**  
**Simple**  $m = D/f$   
**Compound**  
 $m = \frac{v}{u} \frac{D}{f_e}$  Resolving Pow  $R = \frac{1}{\Delta d} = \frac{2\mu \sin \theta}{\lambda}$

**TELESCOPE**  
 $m = -f_o/f_e$   
 $L = f_o + f_e$   
 $R = \frac{1}{\Delta \theta} = \frac{1}{1.22\lambda}$

**DISPERSION**  
 Cauchy's  $\mu = \mu_0 + A/\lambda^2 \quad A > 0$   
 For small A & i  
**mean deviation**  $S_y = (\mu_y - 1)A$   
**Angular dispersion**  $\theta = (\mu_y - \mu_r)A$   
**Dispersive Power**  
 $\omega = \frac{\mu_v - \mu_r}{\mu_y - 1} \approx \frac{\theta}{\delta_y}$

**DISPERSION ONLY**  
 $(\mu_y - 1)A + (\mu'_y - 1)A' = 0$   
**DEVIATION ONLY**  
 $(\mu_v - \mu_r)A = (\mu'_v - \mu'_r)A'$




# HEAT AND TEMP

$F = 32 + \frac{9}{5}C$   
 $K = C + 273.16$   
 Ideal Gas  $\rightarrow PV = nRT$   
 van der Waals  
 $(p + \frac{a}{V^2})(V - b) = nRT$

$L = L_0(1 + \alpha \Delta T)$   
 $A = A_0(1 + 2\alpha \Delta T)$   
 $V = V_0(1 + 3\alpha \Delta T)$   
 THERMAL STRESS  
 $\frac{F}{A} = Y \frac{\Delta L}{L}$

# KINETIC THEORY

EQUIPARTITION OF ENERGY  
 $K = \frac{1}{2}kT$  for each DoF  
 $K = \frac{f}{2}kT$  for  $f$  Degrees of Freedom




Internal Energy  $U = \frac{f}{2}nRT$   $f = 3$  (monatomic);  $5$  (diatomic)

# SPECIFIC HEAT

Specific heat  $s = \frac{Q}{m\Delta T}$   
 Latent heat  $L = Q/m$   
 $C_v = \frac{f}{2}R$   $C_p = C_v + R$   $\gamma = C_p/C_v$   
 $C_v = \frac{n_1 C_{v1} + n_2 C_{v2}}{n_1 + n_2}$   $\gamma = \frac{n_1 C_{p1} + n_2 C_{p2}}{n_1 C_{v1} + n_2 C_{v2}}$

# THERMODYNAMICS


**I<sup>ST</sup> LAW**  $\Delta Q = \Delta U + W$   $W = \int p dV$   
 ADIABATIC  $W = \frac{p_1 V_1 - p_2 V_2}{\gamma - 1}$   
 ISOTHERMAL  $W = nRT \ln(\frac{V_2}{V_1})$   
 ISOBARIC  $W = p(V_2 - V_1)$   
 ADIABATIC:  $\Delta Q = 0$ ;  $PV^\gamma = \text{const}$



# II<sup>ND</sup> LAW ENTROPY $dS = \frac{dQ}{T}$

# HEAT TRANSFER

CONDUCTION  $\frac{\Delta Q}{\Delta t} = -KA \frac{\Delta T}{x}$   
 Thermal Resistance  $= \frac{x}{KA}$



**KIRCHHOFF'S LAW**  $\frac{\text{Emissive Power}}{\text{Absorptive Power}} = \frac{E_{\text{body}}}{a_{\text{body}}} = E_{\text{blackbody}}$   
**WIEN'S DISPLACEMENT**  $\lambda_m T = b$   
**STEFAN-BOLTZMANN**  $\Delta \theta / \Delta t = \sigma e A T^4$   
**NEWTON'S COOLING**  $\frac{dT}{dt} = -bA(T - T_c)$

# ELECTROSTATICS


**COULOMB'S LAW**  $F = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2}$   
 $\vec{E} = \vec{F}/q = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$   
 POTENTIAL (V)  $= \frac{1}{4\pi\epsilon_0} \frac{q}{r}$   
 $PE(U) = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r}$   $\vec{E} = -\frac{dV}{dr}$

**DIAPOLE MOMENT**  $\vec{p} = q\vec{d}$   
**DIAPOLE IN FIELD**  $\vec{p} = \vec{p} \cdot \vec{E}$   $U = -\vec{p} \cdot \vec{E}$   
 $E_r = \frac{1}{4\pi\epsilon_0} \frac{2p \cos \theta}{r^3}$   
 $E_\theta = \frac{1}{4\pi\epsilon_0} \frac{p \sin \theta}{r^3}$   
 $\frac{1}{4\pi\epsilon_0} \frac{p \cos \theta}{r^2} = V(r)$


# GAUSS'S LAW

$\phi = q_{in}/\epsilon_0$  FLUX  $\phi = \oint \vec{E} \cdot d\vec{s}$   
 $E = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \cos \theta$

# UNIFORMLY CHARGED SPHERE



# UNIFORM SHELL



# LINE CHARGE $E = \frac{\lambda}{2\pi\epsilon_0 r}$

$\infty$ -sheet  $E = \frac{\sigma}{2\epsilon_0}$   
 $\vec{E}$  near  $\rightarrow$  CONDUCTING SURFACE  $E = \frac{\sigma}{\epsilon_0}$



# CAPACITORS

$C = q/V$   $C = \epsilon_0 A/d$


**SPHERE**  $C = \frac{2\pi\epsilon_0 L}{\ln(r_2/r_1)}$   
 $C = 4\pi\epsilon_0 \frac{r_1 r_2}{r_2 - r_1}$



PARALLEL  $C_{eq} = C_1 + C_2$  Force b/w plates  $= \frac{Q^2}{2A\epsilon_0}$   
 SERIES  $\frac{1}{C_{eq}} = \frac{1}{C_1} + \frac{1}{C_2}$   $U = \frac{1}{2} CV^2 = \frac{Q^2}{2C} = \frac{1}{2} QV$   
 WITH DIELECTRIC  $C = \epsilon_0 \frac{KA}{d}$


# CURRENT ELECTRICITY

DENSITY  $j = \frac{1}{A} = \sigma E$   
 $V_{drift} = \frac{1}{2m} \frac{eE}{neA}$   
 $R_{wire} = \rho L/A$   $\rho = \frac{1}{\sigma}$   
 $R = R_0(1 + \alpha \Delta T)$   
**OHM'S LAW**  $V = iR$



# KIRCHHOFF'S LAWS

**\* JUNCTION LAW**  $\sum I_i = 0$   
 Sum of all  $i$  towards a node = 0  
**\* LOOP LAW**  $\sum \Delta V = 0$   
 Sum of all  $\Delta V$  in closed loop = 0  
**POWER**  $= i^2 R = V^2/R = iV$



### GALVANOMETER

$i$   $G$   $i_g$   
 $i_g G = (i - i_g) S$   
**Ammeter**  
 $i_g G = (i - i_g) S$   
**Voltmeter**  
 $V_{AB} = i_g(R+G)$

### CAPACITOR

Charging  $q(t) = CV(1 - e^{-t/\tau})$   
 Discharging  $q(t) = q_0 e^{-t/\tau}$   
 Time Constant  $\tau = RC$

### MAGNETISM

$\vec{F}_{LORENTZ} = q\vec{v} \times \vec{B} + q\vec{E}$   
 $qvB = mv^2/r$   
 $T = \frac{2\pi m}{qB}$

### MAGNETIC DIPOLE

$\vec{\mu} = i \text{Area} \hat{z}$   $\vec{\tau} = \vec{\mu} \times \vec{B}$   
 $U = -\vec{\mu} \cdot \vec{B}$   
**HALL EFFECT**  
 $V_w = \frac{Bi}{ned}$

### PELTIER EFFECT

emf  $e = \frac{\Delta H}{\Delta \theta}$

### THOMSON EFFECT


emf  $e = \frac{\Delta H}{\Delta \theta} = \sigma \Delta T$

### FARADAY'S LAW OF ELECTROLYSIS

$m = Zit = \frac{1}{F} Zit$   
 $E = \text{Chem equivalent}$   
 $Z = \text{Electro Chem eq}$   
 $F = 96485 \text{ C/g}$

### SEEBACK EFFECT

$e = aT + \frac{1}{2}bT^2$   
 $T_{neutral} = -a/b$   $T_{inversion} = -2a/b$



### BIOT-SAWART LAW

$d\vec{B} = \frac{\mu_0}{4\pi} \frac{id\vec{l} \times \vec{r}}{r^3}$

$\vec{F} = i \vec{L} \times \vec{B}$

### STRAIGHT CONDUCTOR

$B_{\infty} = \frac{\mu_0 i}{2\pi d}$   
 $B = \frac{\mu_0 i}{4\pi d} [\cos \theta_1 - \cos \theta_2]$

### WIRE

$\frac{dF}{dl} = \frac{\mu_0 i_1 i_2}{2\pi d}$

### AXIS OF RING

$B_p = \frac{\mu_0 i r^2}{2(a^2 + r^2)^{3/2}}$

### CENTER OF ARC

$B = \frac{\mu_0 i \theta}{4\pi r}$   
 $B = \frac{\mu_0 i}{2r}$  (ring)

### SOLENOID

$B = \mu_0 n i$   $n = N/L$

### TOROID

$B = \mu_0 n i$   $n = N/2\pi r$

### AMPERE'S LAW

$\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{in}$

### BAR MAGNET

$B_1 = \frac{\mu_0}{4\pi} \frac{2M}{d^3}$   
 $B_2 = \frac{\mu_0}{4\pi} \frac{M}{d^3}$

### ANGLE OF DIP

$B_h = B \cos \delta$   
 $B_v = B \sin \delta$

### AG

### TANGENT GALVANOMETER

$B_h \tan \theta = \mu_0 n i / 2r$   $i = k \tan \theta$   
**MOVING COIL GALVANOMETER**  
 $n_i AB = k\theta$  ;  $i = \frac{k}{nAB} \theta$

### PERMEABILITY

$\vec{B} = \mu \vec{H}$   
**MAGNETOMETER**  
 $T = 2\pi \sqrt{I/MB_h}$

# ELECTROMAGNETIC INDUCTION

**MAGNETIC FLUX**  $\Phi = \oint \vec{B} \cdot d\vec{s}$  **FARADAY'S LAW**  $e = -\frac{d\Phi}{dt}$   
**LENZ'S LAW:** Induced current produces  $\vec{B}$  that opposes change in  $\Phi$

### SELF INDUCTANCE

$\Phi = Li$   $e = -L \frac{di}{dt}$   
**SOLENOID**  $L = \mu_0 n^2 \pi r^2 l$   
**MUTUAL INDUCTANCE**  
 $\Phi = Mi$  ,  $e = -M \frac{di}{dt}$

### GROWTH

$i = \frac{V}{R} [1 - e^{-t/\tau}]$

### DECAY

$i = i_0 e^{-t/\tau}$   
 Time Const.  $\tau = L/R$   
 ENERGY  $U = \frac{1}{2} Li^2$   
 ENERGY DENSITY OF B-FIELD  
 $u = \frac{U}{V} = \frac{B^2}{2\mu_0}$   
**ROTATING COIL**  $e = NAB\omega \sin \omega t$   
**TRANSFORMER**  $\frac{N_1}{N_2} = \frac{e_1}{e_2}$   
 $C = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$

### ALTERNATING CURRENT

$i = i_0 \sin(\omega t + \phi)$   
 $i_{rms} = i_0 / \sqrt{2}$   
 POWER =  $i_{rms}^2 \cdot R$

### RC-CIRCUIT

$\tan \phi = \frac{1}{\omega CR}$   
 $Z = \sqrt{R^2 + X_C^2}$   
 $X_C = \frac{1}{\omega C}$

### LR-CIRCUIT

$\tan \phi = \frac{\omega L}{R}$   
 $Z = \sqrt{R^2 + X_L^2}$   
 $X_L = \omega L$

### LCR-CIRCUIT

$\tan \phi = \frac{X_C - X_L}{R}$   
 $Z = \sqrt{R^2 + (X_C - X_L)^2}$   
**RESONANCE**  $= \frac{1}{2\pi \sqrt{LC}}$  ( $X_C = X_L$ )  
 $P = e_{rms} i_{rms} \cos \phi$   
 POWER FACTOR

# MODERN PHYSICS

$E = h\nu = hc/\lambda$   $p = h/\lambda = E/c$   $E = mc^2$   
 Ejected photo-electron  $K_{max} = h\nu - \phi$   
**THRESHOLD**  $\nu_0 = \phi/h$   
**STOPPING**  $V_0 = \frac{hc}{e\lambda} - \frac{\phi}{e}$   
 de Broglie  $\lambda = h/p$

### BOHR'S ATOM

**QUANTIZATION OF ANGULAR MOMENTUM**  
 $E_n = -\frac{mZ^2c^4}{8\epsilon_0^2 h^2 n^2} = -\frac{13.6Z^2}{n^2} \text{ eV}$   
 $\gamma_n = \frac{E_n - E_m}{\pi m Z e^2} = \frac{0.529 n^2 A^\circ}{Z}$   $l = \frac{nh}{2\pi}$   
 $E_{TRANSITION} = 13.6 Z^2 (\frac{1}{n^2} - \frac{1}{m^2}) A^\circ$

**HEISENBERG**  $\Delta x \Delta p \geq h/2\pi$   $\Delta E \Delta t \geq h/2\pi$   
**MOSLEY'S LAW**  $\sqrt{\nu} = a(Z - b)$   
**X-RAY DIFFRACTION**  $2d \sin \theta = n\lambda$

# NUCLEUS

$R = R_0 A^{1/3}$  ;  $R_0 = 1.1 \times 10^{-15} \text{ m}$   
**RADIOACTIVE DECAY**  
 $\frac{dN}{dt} = -\lambda N$   $N = N_0 e^{-\lambda t}$   
**HALF LIFE**  $t_{1/2} = 0.693/\lambda$   
**Avg LIFE**  $t_{avg} = 1/\lambda$

### Mass DEFECT

$\Delta m = [Zm_p + (A-Z)m_n] - M$   
**BINDING E**  $= \Delta m \cdot C^2$   
**Q-VALUE**  $Q = U_i - U_f$



# SEMICONDUCTORS

### HALF WAVE RECTIFIER

### TRIODE VALVE

### TRANSISTOR

$I_e = I_b + I_c$   
 $\alpha = \frac{I_c}{I_e}$   $\beta = \frac{I_c}{I_b}$   $\beta = \frac{\alpha}{1-\alpha}$   
 Transconductance  $g_m = \frac{\Delta I_c}{\Delta V_{be}}$

### LOGIC GATES

A	B	AB	A+B	AB	A+B	AB + AB
0	0	0	0	0	0	0
0	1	0	1	0	1	1
1	0	0	1	0	1	1
1	1	1	1	1	0	0

NOW, YOU'RE ONE STEP CLOSER TO YOUR GOAL